

TIME-VARIATIONS OF SOLAR NEUTRINO SIGNALS

P.I. Krastev

*Institute for Advanced Study, Olden Lane, Princeton,
NJ 08540, USA*

The most popular solutions of the solar neutrino problem which assume massive neutrinos predict varying with time signals in the solar neutrino detectors. The variations represent a distinctive feature of these solutions and do not depend on the theoretically predicted total flux of neutrinos from the Sun. The time scale of the variations ranges between an hour (MSW) and several years (neutrino magnetic moment). The amplitude of the variations depends on the neutrino oscillation parameters. Future detectors with high event rates will be able to test these predictions at a new level of precision. The prospects for testing neutrino physics solutions by these detectors will be discussed and the limits on the neutrino oscillation parameters that one should be able to obtain will be presented.

1 Introduction

Solar neutrino experiments are entering the stage of precision measurements. SuperKamiokande¹ and SNO² are going to measure not only the total boron neutrino flux, which alone would be very important, but also the spectrum of recoil electrons and the possible time-variations of the signals. Perhaps the single most important measurement will be the measurement of CC/NC ratio³ in the SNO detector, which might rule out all “new physics” solutions, should the double ratio, $(CC/NC)_{exp}/(CC/NC)_{th}$, of the experimentally measured to the theoretically predicted ratios turn out to be close to one.⁴

The new level of experimental precision requires precise calculations of the expected signals in the future detectors and a careful analysis of the possible implications of all conceivable experimental results. It is well known that astrophysical solutions do not provide a solution of all the solar neutrino problems,⁵ the number of which, instead of diminishing, seems to increase with every new solar neutrino experiment. At present at least two of the three qualitatively different experiments (chlorine, water and gallium) have to be wrong in order for an astrophysical solution to become possible.⁶ Even fitting the experiments with solar neutrino fluxes unconstrained by the complications of solar and nuclear physics, which is taken care of in “state of the art” solar model calculations, but subject only to the luminosity constraint doesn’t solve the problem.⁷ The minimal χ^2 is in an unphysical region with negative beryllium neutrino flux. On the other hand, neutrino oscillations provide an excellent fit to the data without drastically changing solar models or arbitrarily manipulating the experimental results. Still, a solar model independent signature in the

experimentally measured signals, which would be the ultimate proof of lepton number violation, has not been seen yet. Such signatures would be spectral distortions, ratios of different kinds of solar neutrino event rates, preferably in one and the same detector, such as CC/NC, CC/(electron scattering), etc., and time-variations beyond that due to the 7 % energy independent effect from the eccentricity of the earth's orbit. Measurements of time-variations require real-time detectors with sufficiently high event rates and the ability to determine the energy of the incident solar neutrinos. An additional advantage would be the possibility to isolate signals coming from solar neutrinos with a line spectrum, such as ^7Be and/or *pep* neutrinos.

2 Variations due to vacuum oscillations

Variations in this case are due to the eccentricity of the Earth's orbit.⁹ The survival probability is a function which depends strongly on the distance between source and detector. The time-variations are particularly strong for neutrinos with line spectrum, but are large enough to be observed in detectors like SNO and SuperKamiokande. Signals in these detectors are predicted to vary by up to 15 % between minimum and maximum.¹⁰ It is important to use a flux independent normalization of the experimental results, e.g. to divide the average event rate during a month (or a week) by the average event rate during a whole year.¹¹ This will make possible the study of variations independently of the predicted model neutrino flux. The variations are quite sensitive to changes in the detector thresholds, a feature which can be used as an additional check of their nature. Analysis of data obtained during the same time intervals by different detectors, e.g. SNO and SuperKamiokande, can provide further information, thus enhancing the discovery potential of these detectors.

3 Variations due to MSW transitions

Day-night and seasonal variations have been identified a long time ago as particularly revealing features of the MSW effect.^{12,13} The electron neutrinos in the sun are supposedly converted into muon or tau neutrinos inside the sun. During day time these neutrinos are detected on Earth without further modification. At night however, the neutrinos can undergo an additional transition, this time from muon or tau back to the original electron neutrinos, thus enhancing the night-time event rates. This spectacular feature of a solar neutrino signal can in principle be the final, still missing, proof of MSW conversions in the sun. Note that the Kamiokande collaboration has already ruled out a relatively large portion of the $\Delta m^2 - \sin^2 2\theta$ plane, the one where




Figure 1: Asymmetry (day-night)/(day+night) in the SuperKamiokande (a) and SNO (b) detectors.

this effect is particularly strong, by not seeing any sign of variability of their signal.¹⁴ SuperKamiokande is unlikely to do much better even with the superb statistics, because of its sensitivity to the ν_μ -electron scattering, which reduces the amplitude of the day-night variations. On the other hand, SNO has a better chance to probe almost the whole large mixing allowed region and even some portion of the small mixing-angle allowed region. In Fig.1 the asymmetry, (night-day)/(night+day), of the signals within one year between night-time and day-time event rates is plotted in the $\Delta m^2 - \sin^2 2\theta$ plane for the two detectors. Further improvement of the sensitivity of these detectors to the day-night effect can be achieved by analyzing data during shorter time intervals around the winter solstice when the neutrinos pass for a short time through the core of the earth.^{16,17}

4 Conclusions

Variations of solar neutrino signals in future detectors might provide decisive evidence for or against neutrino oscillations.

Acknowledgments

This work has been partially supported by funds from the Institute for Advanced Study.

References

1. M.Takita, in *Frontiers of Neutrino Astrophysics*, edited by Y.Suzuki and K.Nakamura (Universal Academy Press, Tokyo, 1993), p. 147; T.Kajita, *Physics with the SuperKamiokande Detector*, ICRR Report 185-89-2 (1989).
2. Sudbury Neutrino
Observatory Proposal, SNO-87-12 (1987); A.B.McDonald, in Proceedings of the Ninth Lake Louise Winter Institute, edited by A.Astbury *et al.* (World Scientific, Singapore, 1994), p. 1.
3. H.H.Chen, Phys. Rev. Lett. **55**, 1534 (1985).
4. J.N.Bahcall and P.I.Krastev, Phys. Rev. D. **53**, 4211 (1996); J.N.Bahcall and E. Lisi, preprint IASSNS-AST 96/33, (submitted to PRD).
5. J.N.Bahcall and H.Bethe, Phys.Rev. D **47**, 1298 (1993). N.Hata and P.Langacker, Phys. Rev. D **50**, 632 (1993); V. Berezinskii, G. Fiorentini and M. Lissia, Phys.Lett. B **365**, 185 (1996).
6. J.N.Bahcall, Phys.Lett. B **338**, 276 (1994); S.Parke, Phys.Rev.Lett. **74**, 839 (1995); N.Hata and P.Langacker, Phys.Rev. D **52**, 420 (1995).
7. J.N. Bahcall, "How Well Do Standard Solar Models Describe the Results of Solar Neutrino Experiments?", Invited talk at the symposium on The Inconstant Sun, Naples, Italy, March 18, 1996. To be published in *Memorie della Societ *, eds. G. Cauzzi and C. Marmolino.
8. J.N.Bahcall and P.I.Krastev, first paper in ⁴.
9. Ya.Pomeranchuk, cited in next paper; V.N.Gribov and B.M.Pontecorvo, Phys. Lett. B **28**, 493 (1969); J.N.Bahcall and S.C.Frautschi, Phys. Lett. B **29**, 623 (1969); S.L.Glashaw and L.Krauss, Phys.Lett. B **190**, 199 (1987).
10. P.I.Krastev and S.T. Petcov, Phys. Lett. B **299**, 99 (1993).
11. P.I.Krastev and S.T. Petcov, Nucl. Phys. B **449**, 605 (1995).
12. S.P.Mikheyev and A.Yu.Smirnov, in *New and Exotic Phenomena*, Proceeding of the Moriond Workshop, les Arc, savoie, France, 1987, edited by O. Fackler and J. Tran Thanh Van (Editiones Frontieres, Gif-sur Yvette, France, 1987), p./405.
13. A.J.Baltz and J.Weneser, Phys. Rev. D **35**, 528 (1987); Phys.Rev. D **37**, 3364 (1988).
14. By Kamiokande-II Colaboration, Phys. Rev. Lett. **66**, 9 (1991).
15. N.Hata and P.Langacker, Phys.Rev. D **48**, 2937 (1993).
16. A.Baltz and J.Weneser, Phys.Rev.D **50**, 5971 (1994) addendum-ibid. D **51**, 3960 (1995).
17. J.LoSecco, Phys. Rev. D **47**, 2032 (1993).